## **Development of Quantitative Criteria and Rating Scales**

Criteria for evaluating pathogen threats under bioterrorism scenarios, as internal or external introductions (Terms in orange require quantitative scale). Note—need a real zero; use a 1-5 with 3 being the "average"

- 1) Pathogen establishment in the U.S. is possible:
  - a) Infection units (spores, mycelium, sclerotia, etc.) remain viable for a long period of time under natural conditions. 1= less than one day, 2= less than 7 days, 3= less than 30 days, 4 = crop cycle or year, 5 = persists
    - this can be used for all taxa
  - b) There is a natural (wind, vectors, water, etc.) or mechanical (equipment, such as harvesters, sprayers, misters, airplanes crop dusters, etc.) means of **dissemination** within and among growing areas.
  - c) The pathogen has a high infection efficiency.
    - Infection and establishment occurs under a wide range of conditions.
  - d) The pathogen has high reproductive potential in the field
    infection unit/time ; depends on taxa. Ro concept might apply.
  - e) The pathogen has numerous alternative hosts
    - increased risk with asymptomatic **hosts**.
  - f) U.S. **germplasm** is particularly **susceptible** to the pathogen (1=low percentage of available germplasm to 5=100% susceptible).
  - g) The pathogen's U.S. **germplasm** is densely and widely **distributed**. (1=isolated to 5=large acreage of monoculture)
  - h) No effective or economical **control**(s) of the pathogen is **available**.
  - i) Pathogen can survive intercrop periods over a wide range of conditions

## 2) The risk of a particular pathogen can be evaluated by assessment of these characteristic:

- a) The pathogen or its inoculum, or vector is 1=difficult to 5= easy to obtain.
- b) The pathogen or its inoculum, or vector is 1= difficult to 5= easy to grow.
- c) The pathogen or its inoculum, or vector is 1=difficult to 5= easy to handle.
- d) The pathogen or its inoculum, or vector is 1=difficult to 5= easy to transport.
- e) The pathogen or its inoculum, or vector is 1=difficult to 5= easy to deliver.

## 3) A pathway for entry exists

- a) The infested/infected material (IIM) **arrives** at U.S. borders /ports **with frequency**.
- b) The IIM can be **co-mingled** with non-contaminated commodity (-ies), during storage, transport, and /or processing.
- c) The IIM **arrives** at U.S. borders **in volume**, making (sampling/testing/detection) inspection difficult.
- d) The IIM is **distributed** to several locations.
- e) No method for rapid, reliable, and sensitive detection is available on entry.
- f) The pathogen can be **disseminated** by inanimate objects.

## 4) The pathogen has significant social or psychological shock value

a) Pathogen presence can **create uncertainty** or affect markets, whether plants or animals.

- b) The pathogen produces a **toxin** or byproduct, actual or perceived, that **contaminates** or accumulates in **food/feed**.
- c) The pathogen can be **genetically altered** to threaten food/feed security.
- d) The pathogen can **affect natural resources**, **native plants** or ornamentals, and/or urban landscapes.
- 5) Pathogen establishment in the U.S. would have direct economic effects on U.S. farmers, ranchers, or other agricultural producers: [see group 1]
  - a) The commodity (-ies) affected has a high value of production.
  - b) Pathogen presence would adversely **affect** the **market** (raw, processed food/feed, animals).
  - c) Pathogen establishment in the U.S. would **affect** the **economic well-being** of U.S. producers and/or consumers.
- 6) Pathogen establishment in the U.S. or regionally within, would have traderelated economic effects on U.S. farmers, ranchers, or other agricultural producers:
  - a) Pathogen presence would raise unit costs of production [use percentage change in cost] (via yield losses and/or input cost increases) to lessen U.S. comparative advantage in the market.
  - b) The presence of the pathogen in the U.S. would **close off export markets** due to other countries' phytosanitary regulations.
- 7) Public costs of monitoring for, eradicating, or managing the pathogen in the U.S.: Scale of cost is may be direct or inverse to:
  - a) The probability of early detection is low
    - rapid eradication of an introduced pathogen is low.
  - b) The time frame for effective eradication is long.
     1 = days, 2= weeks, 3 = one crop cycle, 4=more than one crop cycle or years 5=unlikely to eradicate
  - c) The costs of the pathogen's eradication would be high.
    - Agronomic versus landscape (leave to economists)
  - d) The costs of monitoring/detecting the pathogen within U.S. borders would be high. (leave to economists)
  - e) The costs of managing the pathogen population would be high. (leave to economists) Dollars/production unit /year
  - f) There is **insufficient knowledge** for producers **to** cost-effectively or successfully **manage** the pathogen.
  - g) Producers and other affected parties do not have the **equipment** and/or **expertise** to deploy **controls** (e.g. chemical, biological, cultural practices).

Threat of quarantine tends to drive disease detection underground