

Rapid Evaluation of Plant Extracts and Essential Oils for Antifungal Activity Against *Botrytis cinerea*

C. L. Wilson, Research Plant Pathologist, J. M. Solar, Chemist, A. El Ghaouth, Visiting Scientist, and M. E. Wisniewski, Plant Physiologist, USDA-ARS Appalachian Fruit Research Station, Kearneysville, WV 25430

ABSTRACT

Wilson, C. L., Solar, J. M., El Ghaouth, A., and Wisniewski, M. E. 1997. Rapid evaluation of plant extracts and essential oils for antifungal activity against *Botrytis cinerea*. Plant Dis. 81:204-210.

A rapid assay to determine antifungal activity in plant extracts and essential oils is described. Wells in microtiter plates were loaded with *Botrytis cinerea* spores and plant extracts or essential oils. Subsequent changes in optical density following spore germination in the wells was measured after 24 h using an automatic microtiter plate reader driven by a software program developed for this purpose. Extracts from 345 plants and 49 essential oils were evaluated for their antifungal activity against *B. cinerea*. Among 345 plant extracts analyzed, 13 showed high levels of antifungal activity, with species of *Allium* and *Capsicum* predominating. Among the 49 essential oils tested, palmarosa (*Cymbopogon martini*), red thyme (*Thymus zygis*), cinnamon leaf (*Cinnamomum zeylanicum*), and clove buds (*Eugenia caryophyllata*) demonstrated the most antifungal activity against *B. cinerea*. The most frequently occurring constituents in essential oils showing high antifungal activity were: *D*-limonene, cineole; β -myrcene; α -pinene, β -pinene; and camphor.

Additional keywords: biocontrol, natural fungicides, volatiles

Among pesticides used to protect crops, fungicides were perceived until recently as relatively safe. A 1986 National Academy of Sciences (NAS) report (8) on pesticide residues on food indicated that fungicides pose more of a carcinogenic risk than insecticides and herbicides together. Therefore, synthetic fungicides are suspect in our food chain, and pressure is increasing to find safer alternatives. Additionally, resistance by pathogens to fungicides has rendered certain fungicides ineffective, creating a need for new ones with alternative modes of action. Present activities to find both natural and synthetic fungicides focus on finding compounds that are safe to humans and the environment.

Plant extracts (4,5) and essential oils (7,9) show antifungal activity against a wide range of fungi. Ark and Thompson (1) showed that garlic extracts contain a potent fungicide. They were able to effectively protect peaches against brown rot (*Monilinia fructicola*) with deodorized

garlic extract preparations. Singh et al. (9) found that essential oils from *Cymbopogon martinii*, *C. oliveri*, and *Trachyspermum ammi* exhibited strong antifungal activity against *Helminthosporium oryzae*.

Davidson and Parish (4) critically reviewed methods used for testing the efficacy of food antimicrobials. Most assays for antimicrobial compounds are labor intensive, time-consuming, and expensive. Microtiter plate methods have proved useful to the pharmaceutical and pesticide industries for rapid screening of bioactive compounds (3). We modified a microtiter plate assay to rapidly detect plant extracts and essential oils that may serve as alternatives to synthetic fungicides for the control of *Botrytis cinerea*.

MATERIALS AND METHODS

Plant extraction method. An extraction method was devised that is simple and does not require added chemicals. Fresh plant material was collected in resealable plastic bags and placed in a freezer for a minimum of 12 h at -20°C . As the plant material was tested, it was withdrawn from the freezer and allowed to thaw for a minimum of 20 min (Fig. 1D). Freezing and thawing fractured the plant cells and caused plant cell fluids to collect in the bag outside the tissue. The plastic bag was tilted so that the fluid collected in one corner. A small hole was cut at the corner, and extracted plant fluids were squeezed through the hole and collected in plastic weight boats (Fig. 1D). The extract was filter-sterilized by passing it through a .22- μm syringe filter (Fig. 1E). A filtered 10% extract solution was prepared by adding 50 ml of the sterile extract with 450 ml of a *B. cinerea* spore suspension (1×10^5 conidia per ml) in sterile malt extract broth.

Preparing the multi-well plates. Fifty ml of the plant extract and spore suspension mixture were pipetted into each well of a row of a 96 multi-well microtitration plate (Flow Laboratories) (Fig. 1F). A nontreated row of a spore suspension was added to each plate as a check, as well as a blank row. Background readings from these checks were subtracted by the software program from wells with fungal growth.

Each well was then sealed with a sheet of dental wax to prevent cross-contamination by volatiles (Fig. 1G). After 24-h incubation of the plates at 24°C , the density of fungal growth in the wells was measured with an SLT Labinstrument Model

Table 1. Plant extracts with the highest persistent antifungal activity^a

Genus	Species	Common name	Family
<i>Adenocalyma</i>	<i>alleaceum</i>	Garlic creeper	<i>Bignoniaceae</i>
<i>Allium</i>	<i>ampeloprasum</i>	Elephant garlic	<i>Liliaceae</i>
<i>Allium</i>	<i>ramosum</i>	Fragrant-flower garlic	<i>Liliaceae</i>
<i>Allium</i>	<i>sativum</i>	Serpent garlic	<i>Liliaceae</i>
<i>Tulbaghia</i>	<i>violacea</i>	Society garlic	<i>Liliaceae</i>
<i>Capsicum</i>	<i>annuum</i>	Pepper 'Chile'	<i>Solanaceae</i>
<i>Capsicum</i>	<i>annuum</i>	Pepper 'Jalapeno'	<i>Solanaceae</i>
<i>Capsicum</i>	<i>annuum</i>	Pepper 'Piquin'	<i>Solanaceae</i>
<i>Capsicum</i>	<i>annuum</i>	Pepper 'Sandia'	<i>Solanaceae</i>
<i>Capsicum</i>	<i>annuum</i>	Pepper 'Sweet'	<i>Solanaceae</i>
<i>Capsicum</i>	<i>annuum</i>	Pepper 'Tepin'	<i>Solanaceae</i>
<i>Capsicum</i>	<i>chinense</i>	Pepper 'Habanero'	<i>Solanaceae</i>
<i>Capsicum</i>	<i>frutescens</i>	Pepper 'Tabasco'	<i>Solanaceae</i>

^a Optical density reading under 40.00 OD. No *Botrytis cinerea* spore germination observed after 48 h at 10% dilution of the crude extract.

Corresponding author: C. L. Wilson
E-mail: cwilson@asrr.arsusda.gov

Accepted for publication 18 November 1996.

Publication no. D-1997-0106-05R

This article is in the public domain and not copyrightable. It may be freely reprinted with customary crediting of the source. The American Phytopathological Society, 1997.

EAR 400 AT microplate reader (Fig. 1A). This instrument was attached to a Wang computer system so that the absorbency-optical density (492-nm filter) could be measured in each well and the difference analyzed.

Operation of microtiter plate reader. A software program was written in BASIC that allowed the operation of the microplate reader from a PC terminal. Once executed, the program checks the filter installed in the reader and displays the

information at the computer terminal. At this point the reader is completely under software control. The program prompts the user for file names and indicates when plates should be changed. The user has the option for a single- or double-plate read-




Fig. 1. Photographs of the assay of plant extracts and essential oils for antifungal activity. (A) Personal computer attached to an SLT Labinstrument Model EAR 400 AT microtiter plate reader. After plates containing extracts and *Botrytis cinerea* spores are incubated for 24 h, they are placed in the tray of the computer-driven reader, which reads each individual well simultaneously for optical density and summarizes the results, which appear on the screen within 1 min. (B) Microtiter plate 48 h after *B. cinerea* spores and extracts have been incubated together at 20°C. Row 3 is a blank with nothing added. Row 8 is a 10% dilution of a *Capsicum* extract. Row 9 is a spore suspension with no plant extract added. (C) Filter paper disks attached to a sheet of dental wax being loaded with essential oils. After the oils are loaded on the disk, the wax sheet is inverted over a microtiter plate and each well is sealed with slight pressure (Fig. 1G). (C₁ insert) Microtiter plate sealed with wax, lid attached and held in place with two large rubber bands. (D) Plant material in resealable bag taken out of the freezer and allowed to thaw. Plant extract is being squeezed through hole cut in corner of bag. (E) Extract being filter sterilized. (F) Dilution of extract and spores being pipetted into wells of microtiter plate. (G) Dental wax being pressed over microtiter wells to seal individual wells.

ing. The single-plate method was used in most instances. When the plant extract is pigmented, the double-plate method was used to distinguish between absorbency from the pigmentation and fungal growth. This method scans a blank plate first—a plate of extracted plant material without spores. The second plate reading contains the plant extract with spores. The software system subtracts the difference in optical density between plates. Both methods analyze 10 treatments with eight replicates and average the data for each treatment. Ten different extract–spore combinations

are placed such that the eight well rows contain the same combination (Fig. 1B). The remaining two rows were used as controls, one with spores and no extract and the other a blank with nothing added. Once scanned, the data were placed in a data file.

In the screen, 10% dilutions of the plant extract were evaluated in the multi-well plates. Density readings ranged from optical density (OD) 470 to 12 on the multiplate reader. Density readings were made after 24 h, and direct observations of spore germination were made with an in-

verted light microscope after 12 and 48 h. A microscope field (40× objective) was chosen at random in each well of the plate, and the percent spore germination was recorded for each of eight wells and averaged.

Plant extracts were divided into four classes based on the OD of the fungal growth in the wells of the microtiter plates and the percent spore germination after 12 and 48 h. The four classes were: (i) OD reading under 40.00 and no *B. cinerea* spore germination observed after 48 h at 10% dilution of the crude extract; (ii) OD reading under 40.00 and less than 10% *B. cinerea* spore germination after 48 h with 10% dilution; (iii) OD reading under 40.00 and less than 50% spore germination after 12 h but more than 80% spore germination after 24 h with a 10% dilution; and (iv) OD reading of over 40.00 and more than 80% germination after 12 h with a 10% dilution. Table 1 represents those extracts described in class 1, Table 2 class 2, Table 3 class 3, and Table 4 class 4.

Evaluation of essential plant oils. Microtiter wells were inoculated with *B. cinerea* spores as described above. Forty-nine essential plant oils were obtained from Aroma Vera (P.O. Box 3609, Culver City, CA 90231). They were extracted by steam distillation or scarification by the manufacturer. To test the volatile fungicidal activity of the oils, 5 µl was placed on small, 8-mm-diameter, filter paper disks attached to the dental wax seal for the multi-well plates (Fig. 1C).

In order to precisely align the filter paper disks, the wax seal was pressed lightly to the top of the multi-well plate so that an imprint of the top of the wells appeared on the wax. Individual filter paper disks were then placed in the center of each well imprint (Fig. 1C). After the essential oils were pipetted onto the disks, the wax seal was placed over the multi-well plate with the filter paper disk down. The wax seal was pressed tightly against the top of the plate to assure that each well was individually sealed. The lid of the plate was then attached, and two large rubber bands were used to hold the lid in place (Fig. 1C₁). Dilutions of the essential oils (50, 25, 12.5, 6.25, 3.13, 1.56, 0.78, and 0.39%) were made with mineral oil, which showed no volatile fungicidal activity.

Spores were germinated at 24°C and observed after 24 and 40 h. The results were recorded as spore germination (+) or no spore germination (–) (Table 4).

Analysis of essential oil components. Chemical constituents of the essential oils were determined with a Hewlett Packard 5890 Series II Gas Chromatograph equipped with a Mass Selective Detector 5971 Series containing a fused silica capillary column, 12 m × 0.02 mm i.d. × 0.33 µm film thickness cross-linked 5% phenyl methyl silicone (HP-1, Hewlett Packard, Wilmington, DE). Conditions employed

Table 2. Plant extracts with intermediate antifungal activity^a

Genus	Species	Common name	Family
<i>Aglaia</i>	<i>odorata</i>	Perfume plant	<i>Meliaceae</i>
<i>Allium</i>	<i>cepa</i>	Potato onion	<i>Liliaceae</i>
<i>Allium</i>	<i>fistulosus</i>	Welsh onion	<i>Liliaceae</i>
<i>Allium</i>	<i>schoenoprasum</i>	Chive	<i>Liliaceae</i>
<i>Allium</i>	<i>tuberosum</i>	Chinese chive	<i>Liliaceae</i>
<i>Capsicum</i>	<i>annuum</i>	'Anaheim'	<i>Solanaceae</i>
<i>Capsicum</i>	<i>annuum</i>	'Ancho'	<i>Solanaceae</i>
<i>Capsicum</i>	<i>annuum</i>	'Big Jim'	<i>Solanaceae</i>
<i>Capsicum</i>	<i>annuum</i>	'Cayenne'	<i>Solanaceae</i>
<i>Capsicum</i>	<i>annuum</i>	'Floral Gem'	<i>Solanaceae</i>
<i>Capsicum</i>	<i>annuum</i>	'Fresno Chile Grande'	<i>Solanaceae</i>
<i>Capsicum</i>	<i>annuum</i>	'Holiday Time'	<i>Solanaceae</i>
<i>Capsicum</i>	<i>annuum</i>	'Hungarian Yellow Wax'	<i>Solanaceae</i>
<i>Capsicum</i>	<i>annuum</i>	'Large Cherry'	<i>Solanaceae</i>
<i>Capsicum</i>	<i>annuum</i>	'Mirasol'	<i>Solanaceae</i>
<i>Capsicum</i>	<i>annuum</i>	'Sante Fe Grande'	<i>Solanaceae</i>
<i>Capsicum</i>	<i>annuum</i>	'Serrano'	<i>Solanaceae</i>
<i>Capsicum</i>	<i>annuum</i>	'Super Chile'	<i>Solanaceae</i>
<i>Capsicum</i>	<i>annuum</i>	'Sweet Roumanian'	<i>Solanaceae</i>
<i>Capsicum</i>	<i>annuum</i>	'Yatsafusa'	<i>Solanaceae</i>
<i>Clematis</i>	<i>paniculata</i>	Sweet autumn clematis	<i>Ranunculaceae</i>
<i>Hesperis</i>	<i>matronalis</i>	Sweet Rocket	<i>Cruciferae</i>
<i>Isatis</i>	<i>tinctoria</i>	Dyer's Woad	<i>Cruciferae</i>
<i>Juglans</i>	<i>nigra</i>	Black walnut	<i>Junglandaceae</i>
<i>Liquidambar</i>	<i>styraciflua</i>	Sweet gum	<i>Hamamelidaceae</i>
<i>Matricaria</i>	sp.	'Barths'	<i>Compositae</i>
<i>Oxalis</i>	<i>europaea</i>	European yellow wood sorrel	<i>Oxalidaceae</i>
<i>Prunus</i>	<i>persica</i>	Peach	<i>Rosaceae</i>
<i>Pyrus</i>	<i>communis</i>	Pear	<i>Rosaceae</i>
<i>Satureja</i>	<i>acinos</i>	Mother of thyme	<i>Labiatae</i>
<i>Taxus</i>	<i>canadensis</i>	American yew	<i>Taxaceae</i>
<i>Taxus</i>	<i>media</i>	Anglo-Japanese yew	<i>Taxaceae</i>

^a Optical density reading under 40.00 OD and less than 10% *Botrytis cinerea* spore germination after 48 h with 10% dilution.

Table 3. Plant extracts with transitory antifungal activity^a

Genus	Species	Common name	Family
<i>Achillea</i>	<i>millefolium</i>	Yarrow	<i>Compositae</i>
<i>Amaranthus</i>	<i>cruentus</i>	Amaranthus	<i>Amaranthaceae</i>
<i>Centranthus</i>	<i>ruber</i>	Red valerian	<i>Valerianaceae</i>
<i>Chrysanthemum</i>	<i>parthenium</i>	Chrysanthemum	<i>Compositae</i>
<i>Chrysanthemum</i>	<i>leucanthemum</i>	Ox-eye daisy	<i>Compositae</i>
<i>Cistus</i>	<i>ladanifer</i>	Laudanum	<i>Cistaceae</i>
<i>Cyperus</i>	<i>esulentus</i>	Nutsedge	<i>Cyperaceae</i>
<i>Gaultheria</i>	<i>procumbens</i>	Wintergreen	<i>Ericaceae</i>
<i>Glycyrrhiza</i>	<i>glabra</i>	Licorice	<i>Leguminosae</i>
<i>Hedera</i>	<i>helix</i>	English ivy	<i>Araliaceae</i>
<i>Linaria</i>	<i>vulgaris</i>	Toadflax	<i>Scrophlariaceae</i>
<i>Pilea</i>	<i>pumila</i>	Clearweed	<i>Urticaceae</i>
<i>Poterium</i>	<i>sanguisorba</i>	Salad burnet	<i>Rosaceae</i>
<i>Prunella</i>	<i>vulgaris</i>	Selfheal	<i>Labiatae</i>
<i>Senecio</i>	<i>obovatus</i>	Squaw-weed	<i>Compositae</i>
<i>Vinca</i>	<i>minor</i>	Periwinkle	<i>Apocynaceae</i>

^a Optical density reading under 40.00 OD and less than 50% spore germination after 12 h but more than 80% spore germination after 24 h with a 10% dilution.

Table 4. Volatile effect of plant essential oils on spore germination of *Botrytis cinerea*: (+) spore germination, (-) no spore germination

Essential oil	Concentration (%)								
	100	50	25	12.5	6.25	3.13	1.56	0.78	0.39
Basil									
(<i>Ocimum basilicum</i>)									
24 h	-								
40 h	+								
Bergamot									
(<i>Citrus bergamia</i>)									
24 h	+								
40 h	+								
Birch									
(<i>Betula nigra</i>)									
24 h	-	-	-	-	+				
40 h	-	-	-	+	+				
Carrot seed									
(<i>Daucus carota</i>)									
24 h	-	-	-	+	+				
40 h	-	+	+	+	+				
Cedar wood									
(<i>Juniperus mexicana</i>)									
24 h	+								
40 h	+								
Chamomile mixta									
(<i>Anthemis mixta</i>)									
24 h	-	-	+	+	+				
40 h	-	+	+	+	+				
Cinnamon leaf									
(<i>Cinnamomum zeylanicum</i>)									
24 h	-	-	-	-	-	-	-	+	+
40 h	-	-	-	-	-	-	-	+	+
Citronella									
(<i>Cymbopogon nardus</i>)									
24 h	-	-	-	-	-	+			
40 h	-	-	-	-	-	+			
Clary sage									
(<i>Salvia sclarea</i>)									
24 h	-	+	+	+	+				
40 h	+	+	+	+	+				
Clove buds									
(<i>Eugenia caryophyllata</i>)									
24 h	-	-	-	-	-	-	-	-	+
40 h	-	-	-	-	-	-	-	+	+
Cypress									
(<i>Cupressus sempervirens</i>)									
24 h	-	+	+	+	+				
40 h	+	+	+	+	+				
Elemi									
(<i>Canarium luzonicum</i>)									
24 h	+								
40 h	+								
Eucalyptus									
(<i>Eucalyptus australiana</i>)									
24 h	-	+	+	+	+				
40 h	+	+	+	+	+				
Fennel									
(<i>Foeniculum vulgare</i>)									
24 h	+								
40 h	+								
Fir									
(<i>Abies balsamea</i>)									
24 h	-	+	+	+	+				
40 h	+	+	+	+	+				
Geranium									
(<i>Pelargonium roseum</i>)									
24 h	-	-	-	-	+				
40 h	-	-	-	+	+				
Grapefruit									
(<i>Citrus paradisi</i>)									
24 h	+								
40 h	+								
Juniper									
(<i>Juniperus communis</i>)									
24 h	+								
40 h	+								

(continued on next page)

Table 4. (continued from preceding page)

Essential oil	Concentration (%)								
	100	50	25	12.5	6.25	3.13	1.56	0.78	0.39
Lavandin (<i>Lavandula fragrans</i>)									
24 h	-	-	-	+	+				
40 h	-	-	+	+	+				
Lavender (<i>Lavandula officinalis</i>)									
24 h	-	-	-	+	+				
40 h	-	+	+	+	+				
Lemon (<i>Citrus limonum</i>)									
24 h	+								
40 h	+								
Lemongrass (<i>Cymbopogon citratus</i>)									
24 h	-	-	-	-	-				
40 h	-	-	-	-	-				+
Lime (<i>Citrus limetta</i>)									
24 h	+								
40 h	+								
Litsea (<i>Litsea cubeba</i>)									
24 h	-	-	-	-	-		+	+	
40 h	-	-	-	-	-		+	+	
Marjoram (<i>Thymus mastichina</i>)									
24 h	-	-	-	+	+				
40 h	-	-	+	+	+				
Mugwort (<i>Artemisia vulgaris</i>)									
24 h	-	-	-	+	+				
40 h	-	-	+	+	+				
Myrtle (<i>Myrtus communis</i>)									
24 h	-	-	+	+	+				
40 h	-	+	+	+	+				
Niaouli (<i>Melaleuca viridiflora</i>)									
24 h	-	-	-	+	+				
40 h	-	+	+	+	+				
Nutmeg (<i>Myristica fragrans</i>)									
24 h	-	+	+	+	+				
40 h	+	+	+	+	+				
Orange (<i>Citrus aurantium</i>)									
24 h	+								
40 h	+								
Oregano (<i>Coridothymus capitata</i>)									
24 h	-	-	-	-	-		-	+	
40 h	-	-	-	-	-		+	+	
Palmarosa (<i>Cymbopogon martini</i>)									
24 h	-	-	-	-	-		-	-	+
40 h	-	-	-	-	-		-	-	+
Patchouly (<i>Pogostemon patchouly</i>)									
24 h	+								
40 h	+								
Peppermint (<i>Mentha piperita</i>)									
24 h	-	-	-	-	+				
40 h	-	-	+	+	+				
Petitgrain (<i>Citrus aurantium</i>)									
24 h	-	-	-	+	+				
40 h	-	+	+	+	+				
Pine (<i>Pinus sylvestris</i>)									
24 h	-	+	+	+	+				
40 h	+	+	+	+	+				

(continued on next page)

Table 4. (continued from preceding page)

Essential oil	Concentration (%)									
	100	50	25	12.5	6.25	3.13	1.56	0.78	0.39	
Rosemary (<i>Rosmarinus officinalis</i>)										
24 h	-	-	+	+						
40 h	-	+	+	+	+					
Rosewood (<i>Aniba roseaodora</i>)										
24 h	-	-	-	-	-					
40 h	-	-	-	-	-	+				
Sage lavandulifolia (<i>Salvia officinalis</i>)										
24 h	-	-	-	+	+					
40 h	-	-	+	+	+					
Savory (<i>Satureja montana</i>)										
24 h	-	-	-	-	-					
40 h	-	-	-	-	-	+				
Peppermint (<i>Mentha piperita</i>)										
24 h	-	-	-	-	-	-	+	+	+	+
40 h	-	-	-	-	-	+	+	+	+	+
Spike (<i>Lavandula spica</i>)										
24 h	-	-	-	+	+					
40 h	-	-	-	+	+					
Spruce (<i>Picea mariana</i>)										
24 h	+									
40 h	+									
Tangerine (<i>Citrus reticulata</i>)										
24 h	+									
40 h	+									
Tarragon (<i>Artemisia dracunculus</i>)										
24 h	+									
40 h	+									
Tea tree (<i>Melaleuca alternifolia</i>)										
24 h	-	-	-	-	+					
40 h	-	-	-	-	+					
Thyme lemon (<i>Thymus hiemalis</i>)										
24 h	-	-	+	+	+					
40 h	-	+	+	+	+					
Thyme red (<i>Thymus zygis</i>)										
24 h	-	-	-	-	-	-	-	-	-	+
40 h	-	-	-	-	-	-	-	-	-	+
Vetiver (<i>Andropogon muricatus</i>)										
24 h	-	+	+	+	+					
40 h	+	+	+	+	+					

were the following: carrier gas, ultra high purity helium (GM Industries, Hagerstown, MD) at 15 psig. Oven temperature program: 45°C (3 min) to 300°C (1 min) at 10°C min⁻¹.

The "as received" essential oils were diluted approximately 200-fold (vol/vol) in HPLC grade methanol (Mallinckrodt Specialty Chemicals, Paris, KY) and injected into the Chromatograph in 1.0-µl aliquots. The major constituent(s) was identified with the aid of a computer-driven algorithm to eliminate possibilities and then by matching the mass spectrum of the analyte with that of a library of electron impact mass spectra.

RESULTS

Plant extracts. Plant extracts showing the greatest antifungal activity were those from species of *Allium* and *Capsicum* (Table 1). A 10% dilution of these extracts registered under 40.00 OD in fungal growth and completely inhibited spore germination of *B. cinerea* after 24 and 48 h.

Another group of plant extracts registered under 40.00 OD in fungal growth and under 10% *B. cinerea* spore germination with a 10% dilution (Table 2). This group was more diverse than those in Table 1 but nevertheless contained predominately species of *Allium* and *Capsicum*.

A more diverse group of plants than

those in Tables 1 and 2 (Table 3) registered under 40.00 OD and showed some transitory fungicidal activity with less than 50% spore germination after 12 h but more than 80% germination after 24 h with a 10% dilution.

Those plant extracts (283 in number) that registered over 40.00 OD on the primary screen with the microtiter plate reader and had more than 80% germination after 12 h with a 10% dilution of the extract were considered to have low or no antifungal activity (data not shown).

Essential oils. Palmarosa (*Cymbopogon martini*) and red thyme (*Thymus zygis*) had essential oils that showed the greatest inhibition of spore germination of *B. cinerea* at

Table 5. Most frequently occurring constituents in the essential oils tested and the plants in which they occurred

Constituent	Plant	Retention time
D-Limonene	<i>Abies balsamea</i>	7.72
	<i>Artemisia vulgaris</i>	7.72
	<i>Citrus aurantium</i>	7.74
	<i>Citrus bergamia</i>	7.70
	<i>Citrus limetta</i>	7.71
	<i>Citrus limonum</i>	7.81
	<i>Citrus paradisi</i>	7.73
	<i>Citrus reticulata</i>	7.70
	<i>Coridothymus capital</i>	7.70
	<i>Cupressus sempervirens</i>	7.73
	<i>Cymbopogon nardus</i>	7.75
	<i>Cymbopogon citratus</i>	7.74
	<i>Litsea cubeba</i>	7.70
	<i>Ocimum basilicum</i>	7.71
	<i>Pelargonium roseum</i>	7.74
Cineole	<i>Pinus sylvestris</i>	7.74
	<i>Rosmarinus officinalis</i>	7.72
	<i>Thymus hiemalis</i>	7.70
	<i>Thymus zygis</i>	7.75
	<i>Artemisia vulgaris</i>	7.72
	<i>Coridothymus capital</i>	7.70
	<i>Eucalyptus australis</i>	7.79
	<i>Lavandula fragrans</i>	7.71
	<i>Lavandula spica</i>	7.78
	<i>Melaleuca alternifolia</i>	7.71
	<i>Myrtus communis</i>	7.74
	<i>Ocimum basilicum</i>	7.70
	<i>Rosmarinus officinalis</i>	7.71
	<i>Salvia officinalis</i>	7.72
	<i>Thymus hiemalis</i>	7.70
β-Myrcene	<i>Thymus mastichia</i>	7.75
	<i>Aniba roseodora</i>	7.71
	<i>Citrus aurantium</i>	7.71
	<i>Cymbopogon martini</i>	7.23
	<i>Cymbopogon nardus</i>	7.21
β-Pinene	<i>Lavandula officinalis</i>	7.21
	<i>Pelargonium roseum</i>	7.21
	<i>Citrus limonum</i>	6.99
α-Pinene	<i>Pinus sylvestris</i>	7.00
	<i>Cupressus sempervirens</i>	6.38
Camphor	<i>Pinus sylvestris</i>	6.40
	<i>Artemisia vulgaris</i>	9.07
	<i>Lavandula spica</i>	9.04
	<i>Salvia lavandulifolia</i>	9.05
	<i>Thymus hiemalis</i>	9.02

the lowest concentration (Table 4). Spore germination was completely inhibited at a 0.78% dilution after 40 h. The next best inhibitor of spore germination was clove buds (*Eugenia caryophyllata*), which showed complete inhibition of spore germination at a 0.78% dilution up to 24 h (Table 4). Another essential oil with notable activity was cinnamon leaf (*Cinnamomum zeylanicum*), which inhibited spore germination at a 1.56% dilution after 40 h. The relative activity of the other volatile oils tested can be found in Table 4.

Essential oil components. The chemical constituents in the essential oils derived by a methanol extraction are presented in Table 5. We are in the process of determining the fungicidal activity of the individual components singly and in combination.

DISCUSSION

Within the large reservoir of natural fungicides that exist in plants and microorganisms, it is reasonable that examples exist that would serve as safe and effective alternatives to synthetic fungicides. Such compounds, if properly formulated and applied, could be used directly or could serve as templates for synthetic analogs. Microbially derived natural compounds have also been proposed as alternatives to synthetic fungicides. Antibiotics such as iturin, produced by *Bacillus subtilis* (6), are effective as fungicides against a number of plant pathogens. However, there is considerable resistance to the use of antibiotics in agriculture. It has been argued that such use will risk the development of resistance in animal pathogens to the antibiotic and thereby diminishes its usefulness in animal disease therapy.

The most plants showing antifungal activity among the 345 plant extracts tested were species of *Allium* or *Capsicum*. Although *Allium sativum* was demonstrated early to have good antifungal activity and to be useful as a postharvest treatment (1), it has not been commercialized. Essential oils have long been recognized as having good fungitoxic compounds (9), but they have not been developed into products for postharvest treatments, since industry finds it easier to patent and protect newly synthesized compounds than natural plant products.

Latent infections are especially difficult to control in harvested commodities, because the pathogen resides in an inactive state within the host tissue. Nonsystemic, synthetic fungicides and biological control agents are ineffective in controlling such infections (10). Natural fungicidal volatiles may be useful in controlling latent infections. Wilson et al. (11) found that a number of the volatiles that emanate from peaches as they ripen are fungicidal. It is possible to fumigate peaches with one of these volatiles (benzaldehyde) and protect them against decay (C. L. Wilson, *unpublished*). Perhaps volatile fungitoxic compounds from plant essential oils could be utilized to control postharvest diseases of fruits and vegetables. With the precipitous

withdrawal of methyl bromide as a fumigant, it may be profitable to explore natural plant volatiles as alternatives.

Natural plant-derived fungicides should provide a wide variety of compounds as alternatives to synthetic fungicides, both as fumigants and as contact pesticides (2). They may also prove valuable as "lead structures" for the development of synthetic compounds. It behooves us to explore more intensely this rich source of fungicides.

ACKNOWLEDGMENTS

We thank Brian E. Otto for his excellent technical assistance and Gary Lightner for writing the software program.

LITERATURE CITED

1. Ark, P. A., and Thompson, J. P. 1959. Control of certain diseases of plants with antibiotics from garlic (*Allium sativum* L.). Plant Dis. Rep. 43:276-282.
2. Cutler, H. G., and Hill, R. A. 1994. Natural fungicides and their delivery systems as alternatives to synthetics. Pages 135-152 in: Biological Control of Postharvest Diseases — Theory and Practice. C. L. Wilson and M. E. Wisniewski, eds. CRC Press, Boca Raton, FL.
3. Daeschel, M. A. 1992. Procedures to detect antimicrobial activities of microorganisms. Pages 57-80 in: Food Biopreservatives of Microbial Origin. B. Ray and M. Daeschel, eds. CRC Press, Boca Raton, FL.
4. Davidson, P. M., and Parish, M. E. 1989. Methods for testing the efficacy of food antimicrobials. Food Technol. 43:148-155.
5. Grange, M., and Ahmed, S. 1988. Handbook of Plants with Pest Control Properties. John Wiley & Sons, New York.
6. Gueldner, R. C., Reilly, C. C., Pusey, P. L., Costello, C. E., and Arendale, R. F. 1988. Isolation and identification of iturins as antifungal peptides in biological control of peach brown rot with *Bacillus subtilis*. J. Agric. Food Chem. 36:366-370.
7. Kurita, N., Makoto, M., Kurane, R., and Takahara, Y. 1981. Antifungal activity of components of essential oils. Agric. Biol. Chem. 45:945-952.
8. Research Council, Board of Agriculture. 1987. Regulating Pesticides in Food — The Delaney Paradox. National Academy Press, Washington, DC.
9. Singh, A. K., Dickshit, A., Sharma, M. L., and Dixit, S. N. 1980. Fungitoxic activity of some essential oils. Econ. Bot. 34:186-190.
10. Wilson, C. L., and El Ghaouth, A. 1993. Multifaceted biological control of postharvest diseases of fruits and vegetables. Pages 181-185 in: Pest Management: Biologically Based Technologies. R. D. Lumsden and J. L. Vaughn, eds. American Chemical Society, Washington, DC.
11. Wilson, C. L., Franklin, J. D., and Otto, B. E. 1987. Fruit volatiles inhibitory to *Monilinia fructicola* and *Botrytis cinerea*. Plant Dis. 71:316-319.