

Increased Expression of the Calmodulin Gene of the Late Blight Fungus *Phytophthora infestans* During Pathogenesis on Potato

Corné M. J. Pieterse, Henk M. Verbakel, Johan Hoek Spaans, Leen C. Davidse, and Francine Govers

Department of Phytopathology, Wageningen Agricultural University, Wageningen, The Netherlands

Received 1 July 1992. Accepted 14 December 1992.

In order to isolate *in planta*-induced genes encoding putative pathogenicity factors of the late blight fungus *Phytophthora infestans*, a genomic library was differentially screened. For the differential hybridization, labeled first-strand cDNA synthesized on mRNA isolated from *P. infestans*-infected potato leaves and on mRNA isolated from the fungus grown *in vitro* were used as probes. This screening resulted in the isolation of the *P. infestans* calmodulin gene. The gene, designated *calA*, contains an open reading frame of 447 base pairs without introns and is unique in the *P. infestans* genome. The predicted amino acid sequence is 89.9–94.6% identical to calmodulins from higher eukaryotes, whereas the identity to calmodulins of higher fungi is significantly less (60.8–85.1%). Expression studies revealed that the *P. infestans calA* gene is constitutively expressed in *in vitro* grown mycelium. However, during pathogenesis on potato the level of *P. infestans* calmodulin mRNA is increased approximately fivefold.

Additional keywords: calcium-binding protein, differential screening, Oomycetes, pathogenicity, potato late blight.

The fungal plant pathogen *Phytophthora infestans* (Mont.) de Bary (Oomycetes), the causal agent of potato late blight, is one of the most devastating diseases of potato (*Solanum tuberosum* (L.)). Infection of potato leaves and tubers by a virulent strain of this hemibiotrophic fungus initially causes a rapidly spreading lesion followed by necrosis, which results in complete destruction of the infected tissue. The specialization of *P. infestans* on solanaceous plants such as potato and tomato implies a mechanism for recognition that most likely requires the exchange of signals between host and pathogen. On the one hand, infection of the host by *P. infestans* induces specific expression of

a particular set of host genes, some of which are thought to be involved in defense (Fritzeimer *et al.* 1987; Taylor *et al.* 1990; Matton and Brisson 1989). On the other hand, growth of the fungus on its host plant can lead to induced expression of genes in the pathogen (Pieterse *et al.* 1992). The products of the *in planta*-induced *P. infestans* genes might be involved in the establishment and maintenance of a compatible interaction between the pathogen and its host plant. Our approach to identify factors that determine the pathogenic abilities of *P. infestans* and factors that are involved in the signaling between host and pathogen is to isolate genes of *P. infestans* that show induced or increased expression during pathogenesis. Identification of the proteins encoded by these genes might elucidate some of the molecular and cellular events involved in plant-pathogen interactions.

To isolate *P. infestans* genes whose expression is specifically induced or increased during colonization of the host plant, we screened a genomic library of *P. infestans* by differential hybridization using labeled cDNA probes synthesized on mRNA isolated from *P. infestans*-infected potato leaves and on mRNA isolated from the fungus grown *in vitro*. This resulted in the isolation of several fungal genes of which the expression is induced during growth of the fungus *in planta*. Characterization of one of the clones led to the identification of an *in planta*-induced gene of *P. infestans* coding for polyubiquitin (Pieterse *et al.* 1991). Here we describe the characterization of another *in planta*-induced gene of *P. infestans* that appeared to encode calmodulin, a calcium-binding protein that has an essential function in various cellular processes.

Calmodulin, which is highly conserved in all eukaryotic organisms, is one of the major intracellular calcium receptors responsible for mediating cellular responses to the calcium fluxes which are generated by a variety of processes. Upon binding of Ca^{2+} , calmodulin undergoes a conformational change, which enables the Ca^{2+} -calmodulin complex to bind to, and in that way to regulate, the activity of a large number of enzymes (Cheung 1980; Roberts *et al.* 1986). It is evident that both the Ca^{2+} and the calmodulin concentration play an essential role in this process. An important class of calmodulin-regulated enzymes are protein kinases whose action is recognized as a primary mechanism for the transduction of extracellular stimuli into intracellular events (Hepler and Wayne 1985; Roberts *et al.* 1986). Thus, through binding of Ca^{2+} , calmodulin can have a regulatory effect on a diverse array of cellular processes.

Corresponding author: C. M. J. Pieterse, Department of Phytopathology, Wageningen Agricultural University, P. O. Box 8025, 6700 EE Wageningen, The Netherlands.

Present address of L. C. Davidse: Biotech Centre Royal Sluis, P. O. Box 22, 1600 AA Enkhuizen, The Netherlands.

The nucleotide sequence data reported will appear in the EMBL, GenBank and DDBJ Nucleotide Sequence Databases under accession number M83535.

MPMI Vol. 6 No. 2, 1993, pp. 164–172

This article is in the public domain and not copyrightable. It may be freely reprinted with customary crediting of the source. The American Phytopathological Society, 1993.

In plants, calmodulin and Ca²⁺ are thought to be involved in the regulation of mitosis, polarized cell growth, and cytoplasmic streaming (Marme and Dieter 1983; Hepler and Wayne 1985). In some fungi, e. g. *Metarhizium anisopliae* (St Leger *et al.* 1989), *Ceratocystis ulmi* (Muthukumar and Nickerson 1984), and *Candida albicans* (Paranjape *et al.* 1990), and in the slime molds *Dictyostelium discoideum* (Lydan and O'Day 1988) and *Physarum polycephalum* (Uyeda and Furya 1986), the Ca²⁺-calmodulin complex has been reported to be involved in growth and differentiation.

In this study we describe the isolation and characterization of the *P. infestans* calmodulin gene (*calA*) and show that the expression level of this gene increases severalfold during pathogenesis. The significance of the *in planta*-induced expression of the *P. infestans calA* gene will be discussed.

RESULTS AND DISCUSSION

An *in planta*-induced gene of *P. infestans* codes for calmodulin.

In order to achieve a 99% probability of having any DNA sequence represented, 80,000 recombinant bacteriophages of the *P. infestans* genomic library were differentially screened. This resulted in the selection of 20 recombinant phage clones containing putative *in planta*-induced genes of the fungus. The number of positive clones does not provide information on the number of *in planta*-induced *P. infestans* genes. One recombinant phage can easily contain a constitutively expressed gene in the proximity of a differentially regulated gene; the latter will not be detectable in the differential screening procedure. In a dot blot experiment, DNA of 10 clones hybridized relatively strongly to the cDNA probe derived from mRNA isolated from infected leaves, whereas hybridization to the cDNA probe synthesized on mRNA isolated from *in vitro* grown mycelium gave relatively weak signals. When the differentially hybridizing DNA fragments from these clones were used as probes on northern blots, it appeared that eight of these 10 clones contain a gene of which the expression is significantly increased during growth of the fungus *in planta* (data not shown). There was no cross-hybridization between these eight clones, indicating that they contain different genes.

In one of the eight positive clones, DHC-C, the coding region of the gene responsible for the differential hybridization was mapped on a single 5.0-kb *SstI* fragment, which was subsequently subcloned into pTZ19U. The other *SstI* fragments from DHC-C did not hybridize to the cDNA probes. In Figure 1, the restriction map of the resulting plasmid, designated pPiCM-S, is depicted. Additional differential hybridization experiments indicated that the gene is located on two linked *PstI* fragments of 1.2 and 1.5 kb, respectively. Dideoxy sequencing of various overlapping subclones within the two *PstI* fragments resulted in the nucleotide sequence shown in Figure 2. In the sequenced region, a single open reading frame (ORF) of 447 nucleotides coding for a protein of 149 amino acids was found. Comparison of the ORF with the nucleotide sequence data library revealed that the gene codes for a

protein that is highly similar to calmodulin, a calcium-binding protein present presumably in all eukaryotes. The deduced amino acid sequence of the gene contains four EF-hand calcium-binding domains (indicated by ★★★ in Fig. 5 below) which is typical for calmodulins. From these data it was concluded that the *in planta*-induced gene present in the recombinant bacteriophage DHC-C encodes calmodulin. Therefore the gene was designated *calA*.

Analysis of the *calA* gene locus.

On a Southern blot containing genomic *P. infestans* DNA digested with various restriction enzymes, a probe derived from the coding region of the *calA* gene (a 342-bp *PvuII/HindIII* fragment from pPiCM-S, see Fig. 1) hybridized to single *BamHI* and *SstI* fragments of 6.0 and 5.0 kb, respectively (Fig. 3). Sequencing data revealed that the coding region of the *calA* gene contains an *EcoRI* and a *PstI* restriction site, and as a result, the *calA* probe hybridizes to two *EcoRI* (2.6 and 17 kb) and two *PstI* fragments (1.2 and 1.5 kb). The length of the hybridizing *SstI* fragment is in agreement with the length of the 5.0-kb *SstI* fragment found in the recombinant bacteriophage DHC-C. The same holds for the two *PstI* fragments and the 2.6-kb *EcoRI* fragment that are present in pPiCM-S (Fig. 1). The sizes of the other hybridizing fragments could not be verified as they overlap the 5.0-kb *SstI* fragment on either one side (17-kb *EcoRI* fragment) or both sides (6.0-kb *BamHI* fragment). From these data it can be concluded that the genome of *P. infestans* contains a single copy of the calmodulin gene. The occurrence of one calmodulin gene per haploid genome is common in most organisms, but in a few cases calmodulin gene families have also been reported (Chung and Swindle 1990; Chien and Dawid 1984; Fischer *et al.* 1988). Each member of these calmodulin gene families may be regulated differently (Fischer *et al.* 1988) but within one organism all calmodulin genes encode identical calmodulin proteins.




Fig. 1. Restriction map and sequencing strategy of the *Phytophthora infestans calA* gene. **A**, Partial restriction map of the genomic λ clone DHC-C. **B**, Restriction map of the 5.0-kb *SstI* insert of pPiCM-S. C = *Clal*, E = *EcoRI*, H = *HindIII*, P = *PstI*, Pv = *PvuII*, S = *SalI*, Ss = *SstI*, and X = *XbaI*. *BamHI* does not cut in the 5-kb insert. The closed bar indicates the approximate location of the coding region of the *calA* gene as determined by differential hybridization experiments. The open bar indicates the coding region of the *calA* gene as determined by dideoxy sequencing. The hatched area in this bar represents the 342-bp *PvuII/HindIII* fragment which was used as probe for northern and Southern blot hybridizations. **C**, Sequencing strategy for *calA*. Arrows indicate the direction and the extent of sequencing.

Further characterization of the *calA* gene.

The nucleotide sequence of the *calA* gene as shown in Figure 2 includes 720 bp upstream of the translation initiation site and 188 bp downstream of the stop codon. The coding region of the *calA* gene is not interrupted by introns. The putative transcription initiation site of the *calA* gene was determined by primer extension using an oligonucleotide complementary to the *calA* sequence at position 9–25 relative to the ATG start codon. The oligonucleotide was annealed to and extended on poly(A)⁺ RNA isolated from 3-day-old *P. infestans* mycelium grown on Henniger synthetic medium. A single primer extension product of 71 nucleotides was found, indicating that the *calA* transcription initiation site is located at the T at position -46 relative to the ATG start codon (Fig. 4). The oligonucleotide used in this primer extension experiment is 82% homologous to the corresponding region in potato calmodulin mRNA (Jena *et al.* 1989). To check whether the oligonucleotide

hybridized to potato calmodulin mRNA, the primer extension experiment was also performed on poly(A)⁺ RNA isolated from the potato-*P. infestans* interaction (potato cv. Ajax, 3 days postinoculation). Again, only one primer extension product was detected that is identical in length to the one produced on poly(A)⁺ RNA isolated from *in vitro* grown mycelium (Fig. 4). Apparently the same *calA* transcription initiation site at position -46 is functional during growth *in vitro* and *in planta*. Potato calmodulin mRNA has a leader sequence of at least 81 nucleotides (Jena *et al.* 1989). Thus the expression level of the potato calmodulin gene is either too low to be detected or the oligonucleotide does not hybridize to potato calmodulin mRNA under the hybridization conditions used.

The transcription initiation site of the *calA* gene is preceded by the sequence TCAT. Interestingly, four other *P. infestans* genes, i.e., the polyubiquitin gene *ubi3R* (Pieterse *et al.* 1991), the actin gene *actA* (Unkles *et al.*




Fig. 2. Nucleotide sequence of the *Phytophthora infestans calA* gene and corresponding amino acid sequence of the encoded calmodulin protein. The transcription start site as determined by primer extension is indicated by a closed arrowhead (▼), the TCATT sequence surrounding the transcription initiation site is underlined with a dotted line. The DNA sequence overlined with the dotted line at positions -51 to -71 represents the sequence motif that is conserved among calmodulin promoter sequences from several organisms. Underlined DNA sequences depict the CAAT and TATA-like motifs. Overlined DNA sequence (AAATAA) in the 3' flanking region of the calmodulin gene represents a putative polyadenylation signal.

1991), and the *in planta*-induced genes *ipiO2A* and *ipiO2B* (C. Pieterse, unpublished) as well as one *Phytophthora megasperma* gene, i.e., the actin gene (Dudler 1990), have a TCATT sequence in which the single transcription initiation site is located. One other *Phytophthora* gene in which the site of transcription initiation has been determined is the actin gene *actB* of *P. infestans* (Unkles *et al.* 1991). However, this gene contains several transcription initiation sites and none of these are located in a sequence motif TCATT. The significance of the conserved TCATT sequence in transcription initiation of *Phytophthora* genes is not clear.

In the promoter region of the *calA* gene, a TATA-like motif (TACAAA) is located 49 nucleotides upstream of the *calA* transcription initiation site at position -95 and a CAAT motif is present at position -167, 121 nucleotides upstream of the transcription initiation site. In the majority of genes identified from higher eukaryotes, the consensus 'core promoter' sequences TATAAA and CAAT are found around 30 bp and 70-90 bp, respectively, upstream of the major transcription initiation site. However, these motifs

are not very common in the promoters of filamentous fungi (Gurr *et al.* 1987). Whether the sequence motifs present in the *calA* gene are functional transcription signals is unknown. Just upstream of the transcription initiation site, between positions -71 and -51, there is a 20 nucleotide sequence motif that is conserved in calmodulin genes of several organisms (Zimmer *et al.* 1988). The conservation of this sequence motif, which is also found in some other unrelated eukaryotic genes (Zimmer *et al.* 1988), ranges between 60 and 80% homology. In the calmodulin genes it is always located within 150 nucleotides upstream of the transcription initiation site. The significance of this motif for transcription or regulation of gene expression is unknown. The 3' untranslated region of the *calA* gene contains a putative polyadenylation signal (AAATAA), 123 nucleotides downstream of the TAA stop codon.




Fig. 3. Southern blot analysis of *Phytophthora infestans* genomic DNA (Pi) digested with *Bam*HI, *Eco*RI, *Pst*I, and *Sst*I and potato genomic DNA (St) digested with *Eco*RI (10 µg/lane). The 342-bp *Pvu*II/*Hind*III fragment from the coding region of the *calA* gene was used as probe. The size of the markers is indicated in kilobases.




Fig. 4. Primer extension analysis of the *calA* transcript. A [γ -³²P]ATP 5'-end-labeled oligonucleotide (5'-GCTCTTCGGTCAGCTGG-3'), complementary to the *calA* sequence at position 9-25, was annealed to 3 µg poly(A)⁺ RNA and extended with M-MLV reverse transcriptase. Primer extension products were electrophoresed on a denaturing 6% (w/v) polyacrylamide gel next to the products of a dideoxy sequencing reaction using the same oligonucleotide as primer and pPiCM-S as template DNA (not shown). The extension products synthesized on poly(A)⁺ RNA isolated from the potato-*Phytophthora infestans* interaction (potato cv. Ajax, 3 days postinoculation) are shown in lane A, the extension products synthesized on poly(A)⁺ RNA isolated from 3-day-old *P. infestans* mycelium grown on Henniger synthetic medium in lane B.

Like most oomycetous genes studied so far, but in contrast to the majority of the known calmodulin genes, the *P. infestans calA* gene has no introns. The few other intron-less calmodulin genes are the calmodulin genes of *Achlya klebsiana* (LeJohn 1989), which is another Oomycete, of *Saccharomyces cerevisiae* (Davis *et al.* 1986) and of *Trypanosoma cruzi* (Chung and Swindle 1990). Most calmodulin genes contain up to five introns, some of which occur at more or less conserved positions in the gene (see references mentioned in the legend of Fig. 5).

Comparison of *P. infestans* calmodulin with calmodulins of other organisms.

The predicted amino acid sequence of the *calA* gene product is highly homologous to calmodulins from other eukaryotic organisms (Fig. 5). The highest identity is found with the calmodulin sequence of the Oomycete *Achlya klebsiana* (96.6%). There are only five amino acid substitutions, which are all located outside the four EF-hand calcium-binding domains. The isoleucine residue at position 109 is unique for *P. infestans* calmodulin. *P. infestans* calmodulin is remarkably similar, 90% identity or more, to calmodulins of several vertebrates, invertebrates, and plants. Also the similarity with calmodulins of lower eukaryotes such as *Trypanosoma cruzi* (91.9%) and *Chlamydomonas reinhardtii* (86.6%), is relatively high. Interest-

ingly, calmodulins of higher fungi such as *Aspergillus nidulans* and especially the yeasts are less homologous to *P. infestans* calmodulin (less than 85% identity). Here, the amino acid substitutions are randomly distributed throughout the sequence. The sequence comparisons suggest that evolutionarily, *P. infestans* is less closely related to higher fungi than to some other eukaryotes. This supports the hypothesis that oomycetous fungi evolved from ancestors other than higher fungi (Barr 1981; Cavalier-Smith 1987; Karlovsky and Prell 1991).

Expression of the *P. infestans* calmodulin gene during pathogenesis on potato.

The *P. infestans calA* gene was isolated from a genomic library by a procedure that was aimed at the selection of *in planta*-induced genes of this fungus. Subsequently, northern blot analyses were performed to confirm that the *calA* gene is indeed a gene that shows induced expression during pathogenesis. When studying gene expression in a plant-fungal interaction by means of northern blot analysis, it is important to take into account the changing ratios of fungal and plant RNA due to the increase of fungal biomass during colonization of the leaves. By using a constitutively expressed fungal gene as probe for the hybridization of northern blots, the increase in the proportion of fungal RNA in the interaction RNA mixtures can

	10	20	30	40	50	60	70	80	
<i>P. infestans</i>	—MADQLTEEQIAEFKFAFSLFDKDG	DTITTKELGTVMRSLGQNPTEAELQDMINEVDADGNGTIDFPEFLTMMARKMKD	TDEE						
<i>A. klebsiana</i>	—G.....			V.....				
Man	—							
Chicken	—							
Eel	—							K.....
<i>D. melanogaster</i>	—							
Potato	—	E.....	C.....			S.A..	Q.....	NL.....	
<i>C. reinhardtii</i>	MAANTE.....		A.....			S.....		ML.....	E.H.D.
<i>T. cruzi</i>	—	SN..S..					Q..S.....	L.....	Q.S.....
<i>A. nidulans</i>	—	S.....	VS.Y.....		Q.....	S.S.....		N.....	
<i>C. albicans</i>	—	EK.S.Q.....		S.K.....		S.S..T.....	VNSD.S.....		A.....
<i>S. pombe</i>	—MTRRN..D.....	R.....	R.Q..N..SN..	V.....	S.A.....			T.....	N.....
<i>S. cerevisiae</i>	—SSN.....	A.....	NN.S.SSS..	A.....	LS.S..VN.LM..	I.V..	HQ.E.S.....	AL.S.QL.SN..	Q.....

	90	100	110	120	130	140	149	% identity
<i>P. infestans</i>	ILEAFKVF	DKDNGF	FISAAELRHIM	TNLGKLTDEE	VDEMIREAD	IDGGQIN	YEEFVKMMMSK*	100.0
<i>A. klebsiana</i>	QG.....		M.....				96.6
Man	.R..R.....	Y.....	V.....			V.....	Q..TA.*	94.6
Chicken	.R..R.....	Y.....	V.....			V.....	Q..TA.*	94.6
Eel	.R..R.....	Y.....	V.....			V.....	Q..TA.*	93.9
<i>D. melanogaster</i>	.R..R.....	Y.....	V.....			V.....	T..T..*	95.9
Potato	LK.....	Q.....	V.....			V.....	R..LA.*	89.9
<i>C. reinhardtii</i>	LR.....		V.....	SE.....	V.....	V.....	R..T.GATDDKDKKHKH*	86.6
<i>T. cruzi</i>	.K..R.....		V.....			V.....		91.9
<i>A. nidulans</i>	.R.....	R.N.....	V..SI.....	D.....	Q...R.D.N..	QL..Q.*		85.1
<i>C. albicans</i>	.A.....	RN.D.K.....	LL.SI.....	S.AD..Q..K..	TNN..E.DIQ..	TLLLA.*		71.6
<i>S. pombe</i>	VR.....	Y.TVE..	T.VL.S..	R.SQ..	AD.....	T...V.....	SRVIS.*	74.5
<i>S. cerevisiae</i>	L.....	N.D.L.....	K.VL.SI.....	A...D.L.VSA..	S.E..	IQQ.AALLA.*		60.8

Fig. 5. Comparison of the predicted amino acid sequence of *Phytophthora infestans* calmodulin with the sequences of calmodulins from: *Achlya klebsiana* (LeJohn 1989), man (Fischer *et al.* 1988), chicken (Putcher *et al.* 1983), *Drosophila melanogaster* (Yamanaka *et al.* 1987), potato (Jena *et al.* 1989), *Chlamydomonas reinhardtii* (Zimmer *et al.* 1988), *Trypanosoma cruzi* (Chung and Swindle 1990), *Aspergillus nidulans* (Rasmussen *et al.* 1990), *Candida albicans* (Saporito and Sypherd 1991), *Schizosaccharomyces pombe* (Takeda and Yamamoto 1987), and *Saccharomyces cerevisiae* (Davis *et al.* 1986). Identical sequences are marked by a dot. The Δ symbol indicates a one-residue gap introduced into the sequence to give optimal alignment with the other proteins. Expected Ca²⁺-binding residues, located in the four EF-hand Ca²⁺-binding domains as proposed by Watterson *et al.* (1980), are indicated by a ★.

be determined. Genes encoding actin and translation elongation factor 1 α (EF-1 α) have been shown to be very useful constitutively expressed marker genes for assessing the expression of differentially regulated genes during plant-pathogen interactions (Mahe *et al.* 1992). In our experiments we used the *P. infestans* actin (*actA*) gene (Unkles *et al.* 1991) as well as the *P. infestans* EF-1 α gene (C. Pieterse, unpublished) as constitutively expressed marker genes. The specificity of the *calA*, *actA*, and EF-1 α probes for *P. infestans* was shown by hybridization of the probes to Southern blots containing genomic *P. infestans* and potato DNA. Under the high-stringency conditions used, the probes hybridized only to *P. infestans* DNA and not to potato DNA (*calA* probe: see Fig. 3; *actA* and EF-1 α probe: data not shown). Also on northern blots the three probes do not hybridize to potato RNA isolated from noninoculated leaves (Fig. 6A).

The expression of the *calA* gene was studied on two different potato cultivars during growth of the fungus *in planta* and during growth of the fungus *in vitro*. Leaves of the partially resistant potato cv. Pimpernel and the moderately susceptible potato cv. Ajax were inoculated with isolate 88069 of *P. infestans*. On potato cv. Ajax, the first symptoms are visible 24 hr after inoculation. They develop from small lesions into completely "water-soaked" areas 3 days postinoculation. In this period the fungus grows and sporulates at the advancing edges of developing

lesions. The centers of the lesions become necrotic and start to decay due to secondary infections by saprophytic microorganisms. Compared to potato cv. Ajax, symptom development on potato cv. Pimpernel was delayed for approximately 1 day. Infected leaves of potato cvs. Ajax and Pimpernel were harvested 1, 2, and 3, and 1, 2, 3, and 4 days postinoculation, respectively. Noninoculated control leaves were sprayed with water and similarly treated to the inoculated leaves. *In vitro* grown mycelium was harvested from nonsporulating cultures (3 day old), moderately sporulating cultures (7 day old) and heavily sporulating cultures (14 day old) which were grown on both Henniger synthetic medium and rich rye-sucrose medium.

On northern blots hybridized with the *calA* probe, a single transcript can be detected in the lanes containing RNA from *in vitro* grown mycelium and in some of the lanes containing RNA isolated from inoculated leaves (Fig. 6). There is no hybridization with RNA isolated from noninoculated leaves. The length of the calmodulin mRNA is about 750 nucleotides, which is in agreement with the length calculated from the nucleotide sequence. In inoculated leaves of potato cv. Ajax as well as those of Pimpernel, the *P. infestans* calmodulin mRNA level increases in time. To be able to normalize calmodulin mRNA levels to a constant amount of fungal RNA, northern blots were deprobed and rehybridized with, successively, the *actA* probe and the EF-1 α probe. During the interaction of *P. infestans*




Fig. 6. A, Autoradiographs of northern blots containing total RNA isolated from noninoculated leaves of potato cvs. Ajax and Pimpernel (n. i. leaves), from colonized leaves of both potato cultivars at different time points postinoculation (days 1-3 and days 1-4, respectively) and from 3-day-old *Phytophthora infestans* mycelium grown on Henniger synthetic medium (*P. infestans*). B, Autoradiographs of northern blots containing RNA isolated from nonsporulating 3-day-old mycelium (3d), from 7-day-old moderately sporulating mycelium (7d), and from 14-day-old heavily sporulating mycelium (14d) grown *in vitro* on both Henniger synthetic medium (HSM) and rich rye-sucrose medium (RSM). Northern blots were hybridized with probes derived from 1) the *P. infestans* calmodulin gene (*calA*), 2) the *P. infestans* actin gene (*actA*), and 3) the *P. infestans* translation elongation factor 1 α gene (EF-1 α).

with potato cv. Ajax, actin and EF-1 α transcripts are first detectable 2 days postinoculation and the amounts increase in time due to the expanding fungal biomass. During the interaction of *P. infestans* with potato cv. Pimpernel both transcripts are first detectable 4 days postinoculation, indicating that the colonization of leaves of the partially resistant potato cv. Pimpernel is significantly slower than the colonization of leaves of the moderately susceptible potato cv. Ajax.

The calmodulin, actin, and EF-1 α mRNA levels during growth of the fungus *in vitro* and *in planta* were quantified by densitometric scanning of the autoradiographs using the Cybertech Image Processing Software, version 1.20 (Cybertech, Berlin). It appeared that the increase in actin mRNA levels due to shifting fungal/plant RNA ratios in the interaction RNA mixtures is comparable to the increase in EF-1 α mRNA levels. Hence, the proportion of fungal RNA in the interaction RNA mixture is quite accurately reflected by the mRNA amounts transcribed from the two constitutively expressed marker genes *actA* and EF-1 α . In addition, the densitometric scanning showed that after normalization to actual fungal RNA levels, the amount of *calA* mRNA in the lanes containing interaction RNA (Ajax, day 2 and 3 and Pimpernel, day 4) is approximately five times higher than in the lanes containing RNA from the *in vitro* grown fungus. Under the *in vitro* growth conditions tested, there is no change in the mRNA levels, indicating that the *calA*, *actA*, and EF-1 α genes are constitutively expressed during growth of the fungus *in vitro*. The higher level of *calA* mRNA during growth of the fungus *in planta* is also demonstrated in the primer extension experiment described above. Equal amounts of primer were annealed to and extended on an identical quantity of poly(A)⁺ RNA isolated from 3-day-old *P. infestans* mycelium grown on Henniger synthetic medium and from the Ajax-*P. infestans* interaction, 3 days postinoculation. The amount of calmodulin mRNA, indirectly visualized by the primer extension products, is severalfold higher in the potato-*P. infestans* interaction compared to *P. infestans* grown *in vitro*. These results strongly suggest that the expression of the *P. infestans calA* gene is significantly increased during colonization of potato. Apparently, growth conditions during pathogenesis directly or indirectly influence the expression level of the *calA* gene.

The role of calmodulin in pathogenesis.

Calmodulin plays an essential role in the regulation of a diverse array of cellular processes. In every eukaryotic cell, a basic level of calmodulin is required for regular cell functions. In this paper we have shown that during colonization of potato by *P. infestans*, the calmodulin mRNA level in the pathogen increases significantly (approximately fivefold).

In several organisms, including fungi, differential expression of calmodulin genes has been observed. In potato, for example, increased levels of calmodulin mRNA are found during tuberization (Jena *et al.* 1989) and in *Drosophila melanogaster* the calmodulin gene is differentially regulated in the various developmental stages (Yamanaka *et al.* 1987). In *Aspergillus nidulans*, calmodulin gene expression changes during the cell cycle (Rasmussen *et al.*

1990). In the oomycetous water molds *Achlya ambisexualis* and *Achlya klebsiana* increased calmodulin gene expression and synthesis are associated with the sporulation process (Suryanarayana and Thomas 1986; LeJohn 1989). To investigate whether the fivefold increase in calmodulin mRNA levels, which is detectable during pathogenesis, is associated with sporulation in the infected tissue, we studied the *calA* gene expression in *in vitro* grown mycelium under sporulating and nonsporulating conditions by northern blot analysis (Fig. 6B). Although the rate of sporulation differs dramatically in the analyzed mycelia, the amount of *calA* mRNA in nonsporulating, moderately sporulating, and heavily sporulating mycelium grown on both synthetic Henniger and rich rye-sucrose medium is similar. This indicates that during growth of the fungus *in vitro*, the expression level of the *calA* gene is not influenced by sporulation nor by the composition of the medium. It can therefore be concluded that induction of calmodulin gene expression during sporulation is not a general phenomenon in Oomycetes.

Pathogenesis is a complex process and in fact very little is known about the properties of *P. infestans* that determine its pathogenic character. Elucidation of the precise reason for increased calmodulin mRNA levels in *P. infestans* during pathogenesis is, at this stage, rather difficult. It has been shown in several organisms that via Ca²⁺-calmodulin complexes, calmodulin is involved in the transduction of extracellular stimuli into intracellular events (Hepler and Wayne 1985; Roberts *et al.* 1986). In view of this, Ca²⁺-calmodulin complexes might be involved in the communication between the two organisms of a plant-microbe interaction. To test the requirement for increased calmodulin levels during pathogenesis, one can think of ways to reduce calmodulin synthesis or calmodulin activity, e.g., by means of anti-sense RNA or via calmodulin inhibitors. Subsequent analyses of the effect of reduced calmodulin synthesis or activity on the pathogenic properties of *P. infestans* can shed more light on the role of calmodulin during pathogenesis. The major drawback of these experiments is the fact that other calmodulin-dependent processes will also be influenced, thus complicating the interpretation of such studies.

The *P. infestans calA* gene was not cloned by the use of heterologous calmodulin probes but was selected via a differential hybridization procedure aimed at the isolation of *in planta*-induced *P. infestans* genes. To our knowledge this is the first observation that a pathogenic plant-microbe interaction is correlated with increased calmodulin mRNA levels in the pathogen. However, at this moment the precise role of calmodulin in the pathogenic process, if any, is still unclear.

MATERIALS AND METHODS

Culturing of *P. infestans* and inoculation of potato leaves.

P. infestans strain 88069 (A1 mating type, race 1.3.4.7), maintained on rye-agar medium containing 2% (w/v) sucrose (Caten and Jinks 1968) was used throughout this study. For inoculations of liquid cultures, zoospores were produced by incubating 2-wk-old rye-agar cultures (in 94

mm petri dishes) for 3 hr at 10° C with a layer of 10 ml of water on top of the mycelium. Liquid cultures were initiated from a zoospore suspension in either Henniger synthetic medium (Henniger 1959) or rich rye-sucrose medium (1 × 10⁵ zoospores per milliliter) and, after incubation at 18° C for 3, 7, and 14 days, mycelium was harvested. For inoculation of plants, sporangia were collected from 2-wk-old rye-agar cultures and concentrated to 5 × 10⁵ sporangia per milliliter. Leaves were inoculated by spraying the axial side of the leaves with the sporangial suspension. After inoculation, leaves were inserted in florist foam oasis and incubated at 18° C and 100% relative humidity under cool fluorescent light (16 hr/day). The partially resistant potato cultivar Pimpinel and the moderately susceptible potato cv. Ajax were used as host plants.

Differential screening, subcloning, and sequencing.

A genomic library of *P. infestans* DNA was constructed in λEMBL3 and screened as described previously (Pieterse *et al.* 1991). For the differential screening, [α -³²P]dATP labeled first-strand cDNA probes with a specific activity of 1–2 × 10⁸ cpm/μg were synthesized on 1) poly(A)⁺ RNA isolated from 3-day-old *P. infestans* mycelium grown on Henniger synthetic medium and 2) poly(A)⁺ RNA isolated from infected leaflets of potato cv. Ajax (3 days post-inoculation) as described by Sargent (1987). Hybridization of filters containing recombinant bacteriophage DNA of the *P. infestans* genomic library was performed at 65° C in hybridization mix containing 5× SSC (750 mM NaCl, 75 mM sodium citrate), 5× Denhardt's solution (0.1% [w/v] Ficoll, 0.1% [w/v] polyvinylpyrrolidone, 0.1% [w/v] BSA [fraction V]), 0.5% (w/v) sodium dodecyl sulphate (SDS), and 100 μg/ml calf thymus DNA. After hybridization, the filters were washed in 2× SSC, 0.5% (w/v) SDS at 65° C, and exposed to Kodak X-AR film.

The location of the *in planta*-induced genes on the DNA inserts present in the selected genomic λ clones was determined by differential hybridization of Southern blots containing digested DNA of the selected λ clones. A single differentially hybridizing 5.0-kb *Sst*I fragment from one of the selected λ clones, i.e. DHC-C, was subcloned in the vector pTZ19U resulting in the plasmid named pPiCM-S. Of pPiCM-S, a detailed restriction map was made and several subclones were constructed. All DNA manipulations were performed essentially according to Sambrook *et al.* (1989). To generate overlapping deletion clones, pPiCM-S was partially digested with *Sau*3AI and electrophoresed on a 0.7% (w/v) agarose gel along with linearized pPiCM-S as marker. Singly cut, linearized DNA was isolated from the gel and digested to completion with *Bam*HI. DNA fragments were then circularized by ligation and transformed to *E. coli* cells. DNA sequencing of overlapping subclones and deletion clones was performed on double-stranded DNA by the dideoxy chain termination method (Sanger *et al.* 1977) using the Multiwell Microtitre Plate Sequencing System (Amersham) and [α -³⁵S]dATP as a label. For the analysis of sequence data and screening of sequence data libraries, the Sequencing Analysis Software Package, version 6.0, of the Genetics Computer Group (GCG) of the University of Wisconsin was used (Devereux *et al.* 1984).

Isolation of RNA and poly(A)⁺ RNA and northern blot analyses.

RNA was isolated from *P. infestans* mycelium, noninoculated potato leaves, and colonized potato leaflets (interaction RNA) using the guanidine hydrochloride RNA extraction method as described by Logemann *et al.* (1987). Poly(A)⁺ RNA was obtained by affinity chromatography on oligo(dT)-cellulose as described by Sambrook *et al.* (1989). For northern blot analysis, 15 μg of total RNA was electrophoresed on denaturing formaldehyde-agarose gels and blotted onto Hybond-N⁺ membranes (Amersham) by capillary transfer as described by the manufacturer. Northern blots were hybridized at 65° C in the hybridization mix as described above.

Isolation of genomic DNA and Southern blot analyses.

Genomic DNA of *P. infestans* and potato was isolated from *in vitro* grown mycelium and from leaves of potato cv. Ajax as has been described previously (Pieterse *et al.* 1991). Genomic DNA was digested with various restriction enzymes and size-separated on a 0.7% (w/v) agarose gel. Following electrophoresis, the DNA was transferred to Hybond-N⁺ membrane (Amersham) by capillary transfer. Hybridization of Southern blots was performed at 65° C in the hybridization mix as described above.

DNA probes.

The calmodulin probe was derived from the coding region of the *P. infestans calA* gene and consists of a 342-bp *Pvu*II/*Hind*III-fragment from pPiCM-S. The 2.8-kb *Pst*I insert from pSTA31 containing the complete *P. infestans actA* gene (Unkles *et al.* 1991) was used as actin probe and the 0.35-kb *Eco*RI/*Xho*I insert from pPi119 which is a partial cDNA clone of the *P. infestans* translation elongation factor 1α (EF-1α) mRNA (C. Pieterse, unpublished), was used as EF-1α probe. DNA probes were labeled by the random primer labeling technique (Feinberg and Vogelstein 1983).

Primer extension of poly(A)⁺ RNA.

The oligonucleotide (5'-GCTCTTCGGTCAGCTGG-3') complementary to the sequence at position 9–25 relative to the ATG start codon of the *calA* gene was 5'-end-labeled with [γ -³²P]ATP using T4 polynucleotide kinase (Promega) according to the manufacturer's instructions. Five nanograms of labeled primer was annealed to 3 μg of poly(A)⁺ RNA and extended using M-MLV reverse transcriptase as described by the manufacturer (Gibco BRL). The primer extension products were analyzed by electrophoresis on a 6% (w/v) polyacrylamide gel and visualized by autoradiography.

ACKNOWLEDGMENTS

We wish to thank Pierre de Wit for continuous support and helpful comments on the manuscript and Maureen Riach for critically reading the manuscript. The genomic clone pSTA31 was kindly provided by Sheila Unkles (University of St. Andrews, St. Andrews, UK). Financial support by the LEB fund is highly appreciated.

LITERATURE CITED

- Barr, D. J. S. 1981. The phylogenetic and taxonomic implications of flagellar rootlet morphology among zoospore fungi. *Biosystems* 14:359-370.
- Caten, C. E., and Jinks, J. L. 1968. Spontaneous variability of single isolates of *Phytophthora infestans*. I. Cultural variation. *Can. J. Bot.* 46:329-347.
- Cavalier-Smith, T. 1987. The origin of fungi and pseudofungi. Pages 339-354 in: *Evolutionary Biology of the Fungi*. A. D. M. Rayner, C. M. Brassier, and D. Moore, eds. Press Syndicate of the University of Cambridge.
- Cheung, W. Y., ed. 1980. *Calcium and Cell Function*, Vol. I. Calmodulin. Academic Press, New York.
- Chien, Y.-H., and Dawid, I. B. 1984. Isolation and characterization of calmodulin genes from *Xenopus laevis*. *Mol. Cell Biol.* 4:507-513.
- Chung, S.-H., and Swindle, J. 1990. Linkage of the calmodulin and ubiquitin loci in *Trypanosoma cruzi*. *Nucleic Acids Res.* 18:4561-4569.
- Davis, T. N., Urdea, M. S., Masiarz, F. R., and Thorner, J. 1986. Isolation of the yeast calmodulin gene: Calmodulin is an essential protein. *Cell* 47:423-431.
- Devereux, J., Haerberli, P., and Smithies, O. 1984. A comprehensive set of sequence analysis programs for the VAX. *Nucleic Acids Res.* 12:387-395.
- Dudler, R. 1990. The single-copy actin gene of *Phytophthora megasperma* encodes a protein considerably diverged from any other known actin. *Plant Mol. Biol.* 14:415-422.
- Feinberg, A. P., and Vogelstein, G. 1983. A technique for radiolabeling DNA restriction endonuclease fragments to high specific activity. *Anal. Biochem.* 132:6-13.
- Fischer, R., Koller, M., Flura, M., Mathews, S., Strehler-Page, M.-A., Krebs, J., Penniston, J. T., Carafoli, E., and Strehler, E. E. 1988. Multiple divergent mRNAs code for a single human calmodulin. *J. Biol. Chem.* 263:17055-17062.
- Fritzemeier, K.-H., Cretin, C., Kombrink, E., Rohwer, F., Taylor, J., Scheel, D., and Hahlbrock, K. 1987. Transient induction of phenylalanine ammonia-lyase and 4-coumarate:CoA ligase mRNAs in potato leaves infected with virulent or avirulent races of *Phytophthora infestans*. *Plant Physiol.* 85:34-41.
- Gurr, S. J., Unkles, S. E., and Kinghorn, J. R. 1987. The structure and organization of nuclear genes of filamentous fungi. Pages 93-139 in: *Gene Structure in Eukaryotic Microbes*. J. R. Kinghorn, ed. IRL Press, Oxford.
- Henniger, H. 1959. Versuche zur kultur verschiedener rassen von *Phytophthora infestans* (Mont.) de By. auf kunstlichen Nahrboden. *Phytopathol. Z.* 34:285.
- Hepler, P. K., and Wayne, R. O. 1985. Calcium and plant development. *Annu. Rev. Plant Physiol.* 36:397-439.
- Jena, P. K., Reddy, A. S. N., and Poovaiah, B. W. 1989. Molecular cloning and sequencing of a cDNA for plant calmodulin: Signal-induced changes in the expression of calmodulin. *Proc. Natl. Acad. Sci. USA* 86:3644-3648.
- Karlovsky, P., and Prell, H. H. 1991. The TRP1 gene of *Phytophthora parasitica* encoding Indole-3-Glycerolphosphate Synthase-N-(5'-Phosphoribosyl) Anthranilate Isomerase. Structure and evolutionary distance from homologous fungal genes. *Gene* 109:161-165.
- Lagace, L., Chandra, T., Woo, S. L. C., and Means, A. R. 1983. Identification of multiple species of calmodulin messenger RNA using a full length complementary DNA. *J. Biol. Chem.* 258:1684-1688.
- LeJohn, H. B. 1989. Structure and expression of fungal calmodulin gene. *J. Biol. Chem.* 264:19366-19372.
- Logemann, J., Schell, J., and Willmitzer, L. 1987. Improved method for the isolation of RNA from plant tissues. *Anal. Biochem.* 163:16-20.
- Lydan, M. A., and O'Day, D. H. 1988. Different developmental functions for calmodulin in *Dictyostelium*: Trifluoperazine and R24571 both inhibit cell and pronuclear fusion but enhance gamete formation. *Exp. Cell Res.* 178:51-63.
- Mahe, A., Grisvard, J., and Dron, M. 1992. Fungal- and plant-specific gene markers to follow the bean anthracnose infection process and normalize a bean chitinase mRNA induction. *Mol. Plant-Microbe Interact.* 5:242-248.
- Marme, D., and Dieter, P. 1983. Role of Ca²⁺ and calmodulin in plants. Pages 263-311 in: *Calcium and Cell Function*, Vol. IV. W. Y. Cheung, ed. Academic Press, New York.
- Matton, D. P., and Brisson, N. 1989. Cloning, expression, and sequence conservation of pathogenesis-related gene transcripts of potato. *Mol. Plant-Microbe Interact.* 2:325-331.
- Muthumukar, G., and Nickerson, K. W. 1984. Ca(II)-calmodulin regulation of fungal dimorphism in *Ceratocystis ulmi*. *J. Bacteriol.* 159:390-392.
- Paranjape, V., Gupta Roy, B., and Datta, A. 1990. Involvement of calcium, calmodulin and protein phosphorylation in morphogenesis of *Candida albicans*. *J. Gen. Microbiol.* 136:2149-2154.
- Pieterse, C. M. J., Risseuw, E. P., and Davide, L. C. 1991. An *in planta* induced gene of *Phytophthora infestans* codes for ubiquitin. *Plant Mol. Biol.* 17:799-811.
- Pieterse, C. M. J., de Wit, P. J. G. M., and Govers, F. 1992, volume 98, supplement 2. Molecular aspects of the potato-*Phytophthora infestans* interaction. *Neth. J. Plant Pathol.*
- Putkey, J. A., Ts'ui, K. F., Tanaka, T., Lagace, L., Stein, J. P., Lai, E. C., and Means, A. R. 1983. Chicken calmodulin genes: A species comparison of cDNA sequences and isolation of a genomic clone. *J. Biol. Chem.* 258:11864-11870.
- Rasmussen, C. D., Means, R. L., Lu, K. P., May, G. S., and Means, A. R. 1990. Characterization and expression of the unique calmodulin gene of *Aspergillus nidulans*. *J. Biol. Chem.* 265:13767-13775.
- Roberts, D. M., Lukas, T. J., and Watterson, D. M. 1986. Structure, function, and mechanism of action of calmodulin. *CRC Crit. Rev. Plant Sci.* 4:311-339.
- Sambrook, J., Fritsch, E. F., and Maniatis, T., eds. 1989. *Molecular Cloning: A Laboratory Manual*, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Sanger, F., Nicklen, S., and Coulson, A. R. 1977. DNA sequencing with chain-terminating inhibitors. *Proc. Natl. Acad. Sci. USA* 74:5463-5467.
- Saporito, S. M., and Sypherd, P. S. 1991. The isolation and characterization of a calmodulin-encoding gene (CMD1) from the dimorphic fungus *Candida albicans*. *Gene* 106:43-49.
- Sargent, T. D. 1987. Isolation of differentially expressed genes. Pages 423-432 in: *Methods in Enzymology*, Vol. 152: Guide to Molecular Cloning Techniques. S. L. Berger, and A. R. Kimmel, eds. Academic Press, San Diego.
- St Leger, R. J., Roberts, D. M., and Staples, R. C. 1989. Calcium- and calmodulin-mediated protein synthesis and protein phosphorylation during germination, growth and protease production by *Metarhizium anisopliae*. *J. Gen. Microbiol.* 135:2141-2154.
- Suryanarayana, K., and Thomas, D. des S. 1986. Asexual spore release from saprolegniaceous watermoulds: Involvement of calmodulin. *J. Gen. Microbiol.* 132:593-598.
- Taylor, J. L., Fritzemeier, K.-H., Hauser, I., Kombrink, E., Rohwer, F., Schroder, M., Strittmatter, G., and Hahlbrock, K. 1990. Structural analysis and activation by fungal infection of a gene encoding a pathogenesis-related protein in potato. *Mol. Plant-Microbe Interact.* 3:72-77.
- Takeda, T., and Yamamoto, M. 1987. Analysis and *in vivo* disruption of the gene coding for calmodulin in *Schizosaccharomyces pombe*. *Proc. Natl. Acad. Sci. USA* 84:3580-3584.
- Unkles, S. E., Moon, R. P., Hawkins, A. R., Duncan, J. M., and Kinghorn, J. R. 1991. Actin in the oomycetous fungus *Phytophthora infestans* is the product of several genes. *Gene* 100:105-112.
- Uyeda, T. Q. P., and Furuya, M. 1986. Effects of low temperature and calcium on microfilament structure in flagellates of *Physarum polycephalum*. *Exp. Cell Res.* 165:461-472.
- Watterson, D. M., Sharief, F., and Vanaman, T. C. 1980. The complete amino acid sequence of the Ca²⁺-dependent modulator protein (calmodulin) of bovine brain. *J. Biol. Chem.* 255:962-975.
- Yamanaka, M. K., Saugstad, J. A., Hanson-Painton, O., McCarthy, B. J., and Tobin, S. L. 1987. Structure and expression of the *Drosophila calmodulin* gene. *Nucleic Acids Res.* 15:3335-3348.
- Zimmer, W. E., Schloss, J. A., Silflow, C. D., Youngblom, J., and Watterson, D. M. 1988. Structural organization, DNA sequence and expression of the calmodulin gene. *J. Biol. Chem.* 263:19370-19383.