

Inheritance of Resistance in Watermelon to Race 1 of *Fusarium oxysporum* f. sp. *niveum*

D. NETZER and CH. WEINTALL, Agricultural Research Organization, Bet Dagan, Israel

ABSTRACT

NETZER, D., and CH. WEINTALL. 1980. Inheritance of resistance in watermelon to race 1 of *Fusarium oxysporum* f. sp. *niveum*. Plant Disease 64:853-854.

The mode of inheritance of resistance to *Fusarium* wilt in watermelon was determined by analyzing the segregation of F₁, F₂, F₃ progenies of crosses involving one susceptible (Mallali) and two resistant (Calhoun Gray, Summit) watermelon cultivars and backcross progenies to the susceptible parental cultivar. The ratios obtained indicate that resistance to race 1 of *Fusarium oxysporum* f. sp. *niveum* is controlled by a dominant gene.

Fusarium wilt, caused by *Fusarium oxysporum* (Schlecht) f. sp. *niveum* (E.F.S.) Snyder & Hans., presents serious problems in growing watermelon in many parts of the world. In Israel, the commercial cultivars as well as those bred recently and introduced during the last decade are susceptible to the local isolates of the *Fusarium* wilt organism (6).

Differences in aggressiveness (*sensu van der Plank*) of isolates of f. sp. *niveum*, without a clear indication of a race situation, have been reported (5). However, Crall (3) reported two races of the fungus in Florida in 1963. Cirulli proved the existence of two races in Italy (2), designated by him as races 0 and 1, differentiated on the basis of a gene-for-gene relationship. Of the three cultivars used by Cirulli, Calhoun Gray was resistant to both races. Paulus et al (8) found this cultivar to provide the highest level of resistance in both field and greenhouse trials. The resistance of Calhoun Gray and Summit to the two races (*sensu* Cirulli) has been corroborated (6). These two cultivars apparently derive their resistance to *Fusarium* wilt from the same source (2).

During a breeding program, we studied the inheritance of resistance in Calhoun Gray and Summit to race 1 of *Fusarium* wilt.

MATERIALS AND METHODS

The inoculum was derived from a culture of *F. oxysporum* f. sp. *niveum* isolated from a wilted watermelon plant in California. This isolate previously (6) was defined as race 1 (*sensu* Cirulli). Details of inoculum preparation and method of inoculation are given elsewhere (7).

Contribution 289-E, 1979 series from the Agricultural Research Organization, The Volcani Center, Bet Dagan, Israel.

Inoculated plants were transplanted into Jiffy pots filled with heat-sterilized sandy loam soil and maintained in a greenhouse with 28 C air and 26 C soil. Calhoun Gray (CG) and Summit (S), both resistant, and the local cultivar Mallali (M), susceptible to the isolate, were used in this study.

Final assessment of wilt symptoms was made on the 15th day after inoculation. After an additional 6 days, the nonwilted plants were transplanted and grown to maturity for either self- or cross-pollinations.

RESULTS

Progenies from self-pollinated M plants were 100% susceptible, whereas progenies from CG or S plants did not

show wilt symptoms. These two latter cultivars were crossed with the susceptible M; none of the F₁ progenies wilted. The segregation observed in the F₂ generation (Table 1) of the crosses M × CG, as well as M × S, suggested simple inheritance (3:1) of the disease reaction, resistance being controlled by one dominant gene. In order to verify this pattern of resistance, the F₁ plants of the crosses CG × M and S × M were backcrossed to the susceptible M. The BC₁ progenies of these crosses gave a 1:1 ratio of resistant to susceptible plants (Table 1).

The mode of inheritance was tested further in F₃ progenies of CG × M, derived from resistant and noninoculated F₂ plants. Fourteen and 10 progenies of such plants, respectively, were obtained (Table 2). Five progenies of the 14 resistant F₂ plants contained resistant plants only, and the nine segregating progenies consisted of a total of 92 resistant and 32 susceptible plants. Of the 10 F₃ progenies of noninoculated F₂ plants, five contained susceptible plants only; two segregating progenies consisted of 34:12 (resistant/susceptible) plants, and three progenies of resistant plants only (Table 2).

Table 1. Segregation in progenies from crosses between resistant (R) cultivars Calhoun Gray and Summit and the susceptible (S) cultivar Mallali after inoculation with race 1^a of *Fusarium oxysporum* f. sp. *niveum*

Parents and crosses	Plants tested (no.)	Expected ratio	Seedlings (no.)		X ²	P
			Resistant	Susceptible		
M	30	all S	0	30
CG	25	all R	25	0
S	15	all R	15	0
F ₁ M × CG	30	all R	30	0
M × S	15	all R	15	0
F ₂ M × CG						
1 ^b	256	3:1	188	68	0.33	0.50-0.70
2 ^c	90	3:1	64	26	0.72	0.30-0.50
Total	346	3:1	252	94	0.86	0.30-0.50
M × S	47	3:1	37	10	0.34	0.50-0.70
BC ₁ (M × CG) × M						
1 ^b	94	1:1	53	41	1.53	0.20-0.30
2 ^d	180	1:1	82	98	1.42	0.20-0.30
Total	274	1:1	135	139	0.05	0.80-0.90
(M × S) × M	26	1:1	15	11	0.61	0.80-0.90

^a*sensu* Cirulli (2).

^bWith Mallali as female parent.

^cWith Calhoun Gray as female parent.

^dWith F₁ plant as female parent.

