

it is much more prevalent and potentially damaging in the cooler climate of the upper Piedmont and mountain regions of the state. While Leath et al (14) suggested that this distribution may be due to reduced or no infection at high temperatures, our data suggest that poor sporulation may also be a factor in restricting the development at higher temperatures. In both this study and that of Leath et al, actual inoculations were done at 26 C; therefore, it was not possible to determine the effects of temperature during the actual infection process. However, the normal nighttime low temperatures during summer on the coastal plain of North Carolina are almost always less than 26 C, so it is doubtful that high temperatures during infection periods are the factor limiting northern leaf blight development in these areas.

LITERATURE CITED

- Adipala, E., Lipps, P. E., and Madden, L. V. 1993. Reaction of maize cultivars from Uganda to *Exserohilum turcicum*. *Phytopathology* 83:217-223.
- Brewster, V. A., Carson, M. L., and Wicks, Z. W., III. 1992. Mapping components of partial resistance to northern leaf blight of maize using reciprocal translocations. *Phytopathology* 82:225-229.
- Carson, M. L. Diversity of types of resistance to northern leaf blight in the BS19 maize synthetic. (Abstr.) *Phytopathology* 82:991.
- Carson, M. L., and Wicks, Z. W., III. 1993. Response of a maize synthetic to S₁ recurrent selection for grain yield in a disease-stress environment. *Maydica* 38:193-199.
- Downs, R. J., and Thomas, J. F. 1991. Phyto-tron procedural manual. N.C. Agric. Res. Serv. Tech. Bull. 244 (revised).
- Gevers, H. O. 1975. A new major gene for resistance to *Helminthosporium turcicum* leaf blight of maize. *Plant Dis. Rep.* 59:296-299.
- Hooker, A. L. 1963. Monogenic resistance in *Zea mays* to *Helminthosporium turcicum*. *Crop Sci.* 3:381-383.
- Hooker, A. L. 1977. Inheritance of chlorotic lesion resistance to *Helminthosporium turcicum* in seedling corn. *Phytopathology* 53:660-662.
- Hooker, A. L. 1977. A second major gene locus in corn for chlorotic-lesion resistance to *Helminthosporium turcicum*. *Crop Sci.* 17:132-135.
- Hooker, A. L. 1980. *Helminthosporium turcicum* as a pathogen of corn. *Rep. Tottori Mycol. Inst. Jpn.* 12:115-125.
- Hooker, A. L. 1981. Resistance to *Helminthosporium turcicum* from *Tripsacum floridanum* incorporated into corn. *Maize Genet. Coop. Newsl.* 55:87-88.
- Hooker, A. L., and Kim, S. K. 1973. Monogenic and multigenic resistance to *Helminthosporium turcicum* in corn. *Plant Dis. Rep.* 57:586-589.
- Hughes, G. R., and Hooker, A. L. 1971. Gene action conditioning resistance to northern leaf blight in maize. *Crop Sci.* 11:180-184.
- Leath, S., Thakur, R. P., and Leonard, K. J. 1990. Variation in expression of monogenic resistance in corn to *Exserohilum turcicum* race 3 under different temperature and light regimes. *Phytopathology* 80:309-313.
- Leonard, K. J., Levy, Y., and Smith, D. R. 1989. Proposed nomenclature for pathogenic races of *Exserohilum turcicum* on corn. *Plant Dis.* 73:776-777.
- Parlevliet, J. E. 1979. Components of resistance that reduce the rate of epidemic development. *Annu. Rev. Phytopathol.* 17:203-222.
- Perkins, J. M., and Pedersen, W. L. 1987. Disease development and yield losses associated with northern leaf blight on corn. *Plant Dis.* 71:940-943.
- Sigulas, K. M., Hill, R. R., Jr., and Ayers, J. E. 1988. Genetic analysis of *Exserohilum turcicum* lesion expansion in corn. *Phytopathology* 78:149-153.
- Thakur, R. P., Leonard, K. J., and Jones, R. K. 1989. Characterization of a new race of *Exserohilum turcicum* virulent on corn with resistance gene HtN. *Plant Dis.* 73:151-155.
- Thakur, R. P., Leonard, K. J., and Leath, S. 1989. Effects of temperature and light on virulence of *Exserohilum turcicum* on corn. *Phytopathology* 79:631-635.
- Ullstrup, A. J. 1970. A comparison of monogenic and polygenic resistance to *Helminthosporium turcicum* in corn. *Phytopathology* 60:1597-1599.
- Ullstrup, A. J., and Miles, S. A. 1957. The effects of some leaf blights of corn on grain yield. *Phytopathology* 47:331-336.

ERRATUM/Volume 78, Number 1, 1994

In the article "Contribution of Four Races of *Xanthomonas campestris* pv. *vesicatoria* to Bacterial Spot in Barbados" by Leonard W. O'Garro and Simone Tudor on pages 88-90, the subhead over column 5 in Table 2 should read: Pepper group race 1 only.